

Synthesis, characterization, and quantum chemical studies of new chiral acyclic (O,N,Te) type Schiff base, (*R*)-1-(((1-((4-methoxyphenyl)tellanyl)propan-2-yl)imino)methyl)naphthalen-2-ol

H. R. Rajegowda^{1,4}, Padala Ashok², Chethan B. S.³ and P. Raghavendra Kumar^{4*}

{rajas.gowda@gmail.com¹, bschethan1995@gmail.com³ and raghukp1@gmail.com⁴}

¹Department of Chemistry, Acharya Institute of Graduate Studies, Soladevanahalli – 560107, Bengaluru, Karnataka, India

²Department of Physics, Vignana Bharathi Institute of Technology, Hyderabad, Telangana-501301, India

³Department of Physics, Sri Siddhartha Institute of Technology, Tumakuru 572105. Karnataka, India

⁴Department of Chemistry, University College of Science, Tumkur University, Tumakuru – 572103, Karnataka, India

Abstract. The reaction of the chiral telluro-amine, (*R*)-2 with 2-hydroxy-1-naphthaldehyde has produced a new chiral telluro Schiff's base of (O,N,Te) type, (*R*)-1-(((1-((4-methoxyphenyl)tellanyl)propan-2-yl)imino)methyl)naphthalen-2-ol (*R*)-3. The SOR, CHN and spectroscopic analysis were used to characterize the structure of (*R*)-3. The different intermolecular interactions were shown by the fingerprint plots and Hirshfeld surface analysis. To describe the optimal structure, FMO, and reactive characteristics, DFT studies were carried out. Additionally, the RDG-based isosurface analysis conducted to investigate the type and nature of interactions.

Keywords: Schiff's base, chiral, tellurium, DFT, QTAIM

1 Introduction

As chiral ligands in transition metal-mediated asymmetric organic transformations [12–14], and in biochemistry [6–11], schiff bases of the organochalcogen (organosulphur, organoselenium, and organotellurium) compounds [1–5] have attracted a lot of attention because of their potential uses. A variety of hybrid organosulfur/selenium/tellurium ligands, as well nitrogen/oxygen ligand chemistry has been thoroughly researched over the past three decades [15–18]. Additionally, organotellurium compounds shown superior antioxidant activity in comparison to their comparable selenium and sulphur compounds [19,20]. In contrast to chiral organo-seleno- and -thioether Schiff bases compounds [21–25], the known

chiral tellurated Schiff bases compounds are extremely rare and have been studied sparsely in biology [6–11, 15–20] and asymmetric organic synthesis. By the reaction of tellurated aldehydes with various chiral amines, the corresponding chiral telluro Schiff baseligands were reported [26–28] but this method is limited due to non commercial availability of the tellurated aldehydes and their synthesis is as not easy as like other.

Chemical and biological research has benefited immensely from the development of computational tools in current years, which have made to facilitate in investigation of basic physical and chemical characteristics and comprehend the nature of noncovalent interactions. Theoretical and experimental studies further improve the structures and characteristics of substances [29]. There are some known chiral tellurium-ether ligands, and their computational behavior has been insufficiently explored.

As a result, we have described the new chiral tellurium based Schiff base (*R*)-**3**, which was synthesized by utilizing chiral tellurated amine. (*R*)-**3** was characterized by FT-IR spectroscopy, elemental analysis, polarimetry, ¹H-NMR, and ¹³C{¹H}-NMR. Furthermore, the QTAIM analysis and DFT computations were carried out.

2 Experimental

2.1 Materials and Methods

Merck Ind. Ltd. provided the tellurium, anisole, (*R*)-2-Amino-1-propanol, 2-Hydroxy-1-naphthaldehyde, and sodium borohydride (NaBH₄), which were utilized exactly as supplied. Standard procedures were used to purify and dry the solvents as needed [30]. Merck 60 F₂₅₄ silica gel pre-coated aluminum TLC plates were used to monitor the reaction and the results were observed using UV light, KMnO₄ solution, or ninhydrine stain.

As described in the literature, the precursor compounds bis (4-methoxyphenyl) ditelluride (Ar₂Te₂) [31,32] and chiral telluro-amine, (*R*)-**2** [28] were made. Melting points of the compounds measured, and the results were reported uncorrected. A cell with a path length of 100 mm was used to measure the specific optical rotation (SOR) using a Rudolph Autopol-I automatic polarimeter. ¹H-NMR and ¹³C{¹H}-NMR spectras were recorded by using tetramethylsilane (TMS) as an internal standard (0 ppm) and the values are expressed in δppm.

2.2 Synthesis of telluro Schiff base, (*R*)-**3**

At room temperature, telluroamine, (*R*)-**2** (1 mmol) dissolved in absolute ethanol, which is added to 2-hydroxy naphthaldehyde solution (1 mmol) dissolved in absolute ethanol with stirring, continued for 3 hrs. After employing a rotary evaporator pale yellow solid resulted of (*R*)-**3**. (Yield: 0.38 g, 85%); M.P.: 77–79 °C; $[\alpha]_D^{26} = -178^\circ$; Anal. Calcd. (found) for C₁₇H₁₈N₂OS: C 42.91 (42.78), H 3.43 (3.34), N 2.38 (2.25); FTIR (ATR) ν , cm⁻¹: 3436 (O-H), 2919 (C-H), 1629 (C=N), 690 (ArC-Te), 606 (alkylC-Te); ¹H-NMR (Solvent:DMSO-d₆, 400 MHz) δ ppm: 1.343–1.359 (doublet, *J*=6.4Hz, 3H, methyl), 2.691–2.742 (doublet of doublet, *J*=7.2Hz, 1H, -CH₂-), 2.795–2.841 (doublet of doublet, *J*=7.6Hz, 1H, -CH₂-), 3.811 (singlet, 3H, OCH₃), 3.846–3.893 (sextet, 1H, CH-N), 6.753–6.776 (doublet, *J*=9.2Hz, 2H, ArH, *meta* to Te), 7.130–7.263 (multiplet, 3H, ArH), 7.419–7.456 (doublet, 1H, ArH), 7.623–7.642 (doublet, 1H, ArH), 7.712–7.736 (doublet, 1H, ArH), 8.062–8.083 (doublet, *J*=8.4Hz, 2H,

ArH, *ortho* to Te), 9.080-9.103 (doublet, $J=9.2\text{Hz}$, 1H, -CH=N), 13.154 (singlet, 1H, OH); $^{13}\text{C}\{^1\text{H}\}$ -NMR (Solvent:DMSO- d_6 , 100.64 MHz) δ : 21.895 (CH_3), 27.343 (C-2), 31.605 (OCH_3), 58.428 (C-1), 119.137 (C-8), 122.795 (C-16), 125.179 (C-10), 125.947 (C-5), 127.027 (C-14), 128.321 (C-15), 128.808 (C-13), 129.314 (C-12), 134.509 (C-11), 137.233 (C-17), 139.782 (C-4), 157.442 (C-9), 159.282 (C-6), 175.916 (C-7).


2.3 Quantum computational studies

The LANL2DZ and B3LYP functional basis set are used in DFT to optimize the geometry of (R)-**3**. The absence of imaginary frequencies showed that the optimized structure corresponded to the true energy minima configuration. To investigate the electronic-properties, global-reactivity parameters, and electrostatic-potential-surface, the FMO and MEP map were plotted respectively and examined at the same theoretical level. All calculations were done by using the Gaussian 16, and the Gaussview 5.0 tool was utilized to show the DFT findings [33]. Furthermore, the reduced density gradient (RDG) based isosurface was developed. The Bader's QTAIM analysis were conducted in order to understand the various types of interactions displayed by (R)-**3**. The Multiwfn software [34] was utilized to perform the QTAIM analysis, and Visual Molecular Dynamics (VMD) tool [35] was used to construct the isosurface plot.

3 Results and discussion

3.1 Synthesis

Chiral telluro Schiff base, (R)-**3** has been synthesised through the condensation reaction of telluro amine, (R)-**2** with 2-hydroxynaphthaldehyde as mentioned in **Scheme 1**. Schiff base, (R)-**3** was found insoluble in n-hexane, heptane, THF, benzene, toluene, and diethyl ether but freely soluble in CH_3CN , DMF, CHCl_3 , CH_2Cl_2 , CH_3OH and DMSO.



Scheme 1 Synthesis of (*R*)-**3**

3.2 Spectroscopy

The compound (*R*)-**3** was characterized by, SOR, CHN analysis, IR, proton and carbon-13 NMR spectroscopy.

As per the FT-IR data of (*R*)-**3**, the O-H and >C=N- stretching vibrational frequencies were appeared at 3436 and 1629 cm⁻¹. The ArC-Te and RC-Te stretching frequencies were appeared at 690 cm⁻¹ and 606 cm⁻¹. The appeared frequencies are found characteristic with the respective functional groups as reported in the literature. [36]




Fig. 1 FT-IR spectrum of (R)-3

In the proton Nuclear Magnetic Resonance spectrum of (R)-3 (as shown in Fig. 2), At the low field region, the aromatic OH proton was appeared, at 14.3 δ ppm it was found as broad singlet and highly deshielded due to hydrogen bonding with imine nitrogen. The proton of the imine (CH=N) was acquired. at δ , 9.1 ppm. The asymmetric CH proton shown a broad quartet at δ , 3.85-3.89 ppm due to coupling with three protons of CH₃ and CH₃ protons formed a doublet at δ , 1.30-1.35 ppm due to coupling with one proton of CH. At δ , 2.69-2.74 and 2.79-2.84 ppm, a series of doublets was produced by diastereoid CH₂Te protons that are chemically non-equivalent. At their distinctive locations (δ ppm), the additional aromatic proton signals were visible [36,37].




Fig. 2 ¹H NMR spectrum of (R)-3

In the carbon-13 Nuclear Magnetic Resonance spectrum of (*R*)-**3** (as shown in **Fig. 3**) the CH₂Te-C signal was obtained at 21.89 δ ppm. The CH -C gave peak at 58.43 δ ppm and imine carbon (C=N) was obtained at δ , 175.92 ppm and was found more deshielded. The Aromatic Carbon-OH peak was obtained at 159.28 δ ppm. The ArC-Te carbon signal was appeared at δ , 106.31 ppm. At their distinctive locations (δ ppm), the other carbon signals were visible [36,37].




Fig. 3 $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of (*R*)-**3**

3.3 Quantum computational studies

The Schiff base, (*R*)-**3** optimized structure is shown in the **Fig. 4** and the HOMO and LUMO was calculated [38]. The HOMO-LUMO gap of (*R*)-**3** was obtained as 3.465 eV indicating the charge transfer occurring within the (*R*)-**3** molecule. The electrophilic behaviour of (*R*)-**3** was revealed by the higher electrophilicity index (3.665 eV) and lower chemical potential value (-3.564 eV). The HOMO and LUMO energies provide the valuable insights into several key parameters including the electronegativity (χ), energy gap (ΔE_g), electron affinity (EA), ionization energy (IE), chemical potential (μ), global softness (s), electrophilicity (ω) and global hardness (η) of (*R*)-**3**, as tabulated in Table 1. An essential measure of a compound's stability and reactivity is its energy gap; a higher gap indicates more molecular stability and aromaticity as well as less chemical reactivity. The frontier molecular orbital distribution of (*R*)-**3** is as shown in **Fig. 5**. It is disclosed that the LUMO orbital is confined around the 2-hydroxy naphthalene moiety, while the HOMO is dispersed over the tellurium atom.




Fig. 4 Optimized structure of (R)-3

Table 1: The global quantum chemical parameters of (R)-3

Parameters	Values
E_{LUMO} (eV)	-1.831
E_{HOMO} (eV)	-5.296
ΔE_g (eV)	3.465
IE (eV)	5.296
EA (eV)	1.831
χ (eV)	3.564
η (eV)	1.732
s (eV) ⁻¹	0.577
ω (eV)	3.665
μ (eV)	-3.564

The optimal structure of (R)-3 was obtained using the QTAIM analysis, and the VMD program was used to create the RDG-based isosurface plot. The isosurface plot of (R)-3 is shown in the **Fig. 6** which reveals the existence and nature of various interactions. The varied color codes in the plot reflect the type of interaction. The existence of interactions of the van der Waals kind is shown by the green regions that are seen between the neighbouring atoms. Interestingly, the presence of the ring coloured region was observed between the carbon and the tellurium atom connected to it. All the rings of aromatic benzene are found to be associated with the red colored region in the center indicating the presence of steric repulsion.




Fig. 5. Frontier molecular orbital distribution of (*R*)-3




Fig. 6 RDG based isosurface of (*R*)-3

Conclusion

A new chiral tellurium based Schiff base, (*R*)-3 was synthesized by the reaction of telluro-amine, and (*R*)-2 and 2-hydroxy-1-naphthaldehyde. The synthesized Schiff's base is

characterized by SOR, elemental and spectroscopic analysis. The geometrical coordinates of (*R*)-**3** was optimized using DFT and the orbital energy gap between HOMO-LUMO of (*R*)-**3** is found to be 3.465 eV. Further, the electrophilic behaviour of (*R*)-**3** was revealed by the higher electrophilicity index (3.665 eV) and lower chemical potential value (-3.564 eV). The HOMO orbital is found to be distributed over the telluriu atom, whereas the LUMO orbital is found to be localized over the 2-hydroxy naphthalene moiety. The isosurface analysis revealed the nature and type of interactions exhibited by (*R*)-**3**.

Acknowledgement

P Raghavendra Kumar acknowledges the financial assistance provided by DST-SERB, New Delhi, India, for the major research project (DST/SR/S1/IC-76/2010(G)).

References

- [1] Schneider, A., Rodrigues, O. E., Paixão, M. W., Appelt, H. R., Braga, A. L., & Wessjohann, L. A.: Stereoselective synthesis of Boc-protected L-seleno- and tellurolanthionine, L-seleno- and tellurocystine and derivatives. *Tetrahedron letters*. 47(6) pp. 1019-1021 (2006)
- [2] Braga, A. L., Galetto, F. Z., Taube, P. S., Paixao, M. W., Silveira, C. C., Singh, D., & Vargas, F.: Mild and efficient one-pot synthesis of chiral β -chalcogen amides via 2-oxazoline ring-opening reaction mediated by indium metal. *Journal of Organometallic Chemistry*. 693(24) pp. 3563-3566 (2008)
- [3] Vargas, F., & Comasseto, J. V.: Practical synthesis of chiral β -telluro amines by ring-opening reaction of aziridines. *Journal of Organometallic Chemistry*. 694(1) pp. 122-126 (2009)
- [4] Affeldt, R. F., Braga, H. C., Baldassari, L. L., & Luedtke, D. S.: Synthesis of selenium-linked neoglycoconjugates and pseudodisaccharides. *Tetrahedron*. 68(51) pp. 10470-10475 (2012)
- [5] Kumar, A., Agarwal, M., Singh, A. K., & Butcher, R. J.: Palladium (II), platinum (II), ruthenium (II) and mercury (II) complexes of potentially tridentate Schiff base ligands of (E= N, O) type (E=S, Se, Te): Synthesis, crystal structures and applications in Heck and Suzuki coupling reactions. *Inorganica Chimica Acta*. 362(9) pp. 3208-3218 (2009)
- [6] Ba, L. A., Döring, M., Jamier, V., & Jacob, C.: Tellurium: an element with great biological potency and potential. *Organic & biomolecular chemistry*. 8(19) pp. 4203-4216 (2010)
- [7] Savegnago, L., Vieira, A. I., Seus, N., Goldani, B. S., Castro, M. R., Lenardão, E. J., & Alves, D.: Synthesis and antioxidant properties of novel quinoline-chalcogenium compounds. *Tetrahedron Letters*. 54(1) pp. 40-44 (2013)
- [8] Andreadou, I., Menge, W. M., Commandeur, J. N., Worthington, E. A., & Vermeulen, N. P.: Synthesis of novel Se-substituted selenocysteine derivatives as potential kidney selective prodrugs of biologically active selenol compounds: evaluation of kinetics of β -elimination reactions in rat renal cytosol. *Journal of medicinal chemistry*. 39(10) pp. 2040-2046 (1996)
- [9] Nogueira, C. W., Zeni, G., & Rocha, J. B.: Organoselenium and organotellurium compounds: toxicology and pharmacology. *Chemical reviews*. 104(12) pp. 6255-6286 (2004)
- [10] Wollenhaupt, S. G., Soares, A. T., Salgueiro, W. G., NoreMBERG, S., Reis, G., Viana, C., & Avila, D. S.: Seleno- and telluro-xylofuranosides attenuate Mn-induced toxicity in *C. elegans* via the DAF-16/FOXO pathway. *Food and Chemical Toxicology*. 64 pp. 192-199 (2014)

- [11] Rajegowda, H. R., Raghavendra, K. P., & Udaya Bhanu, S. D.: Synthesis, Characterization and in vitro Antioxidant Activity of New Chiral N-boc Organotellurium Compounds, $(\text{CH}_3)_3\text{OC}(\text{O})\text{NHCH}(\text{R})\text{C}(\text{O})\text{NHCH}_2\text{-CH}_2\text{Te-C}_6\text{H}_4\text{-4-OCH}_3$, Containing Carbamate and Peptide Groups. *J. Braz. Chem. Soc.* 27(7) pp. 1157-1164 (2016)
- [12] Pamies, O., Margalef, J., Canellas, S., James, J., Judge, E., Guiry, P. J., ... & Dieguez, M.: Recent advances in enantioselective Pd-catalyzed allylic substitution: from design to applications. *Chemical reviews.* 121(8) pp. 4373-4505 (2021)
- [13] Schwab, R. S., Soares, L. C., Dornelles, L., Rodrigues, O. E., Paixao, M. W., Godoi, M., & Braga, A. L.: Chiral chalcogen peptides as ligands for the catalytic enantioselective aryl transfer reaction to aldehydes. pp. 3574-3578 (2010)
- [14] Sehnem, J. A., Milani, P., Nascimento, V., Andrade, L. H., Dorneles, L., & Braga, A. L.: Synthesis of new fluororous modular chiral ligand derivatives from amino alcohols and application in enantioselective carbon-carbon bond-forming alkylation reactions. *Tetrahedron: Asymmetry.* 21(8) pp. 997-1003 (2010)
- [15] Doering, M., Ba, L. A., Lilienthal, N., Nicco, C., Scherer, C., Abbas, M., & Jacob, C.: Synthesis and selective anticancer activity of organochalcogen based redox catalysts. *Journal of medicinal chemistry.* 53(19) pp. 6954-6963 (2010)
- [16] Kumar P, R., Singh, A. K., Butcher, R. J., Sharma, P., & Toscano, R. A.: Tellurated Schiff Bases Formed from $\{2\text{-}[(4\text{-Methoxyphenyl})\text{telluro}]ethyl\}$ amine and Bis(2-aminoethyl)Telluride with o-Hydroxyacetophenone: Synthesis and Complexation Reactions with HgII, PdII and RuII- Crystal Structures of the Ligands, $[\text{Ru}(\text{p-cymene})\text{Cl}\{\text{H}_2\text{NCH}_2\text{CH}_2\text{TeC}_6\text{H}_4\text{-4-OCH}_3\}]\text{Cl}\cdot\text{H}_2\text{O}$ and $[\text{RuCl}\{4\text{-MeOC}_6\text{H}_4\text{TeCH}_2\text{CH}_2\text{NHCH}(\text{CH}_3)\text{C}_6\text{H}_4\text{-2-O-}\}]\text{Cl}$. *European Journal of Inorganic Chemistry.* 5 pp. 1107-1114 (2004)
- [17] Pandey, R. K., Singh, A. K., Drake, J. E., Hursthouse, M. B., & Light, M. E.: First structurally characterized complex of an acyclic tellurated Schiff base 4-MeOC₆H₄TeCH₂CH₂N=C(CH₃)C₆H₄-2-OH ((3)-H-1) having metal-tellurium bond; synthesis and crystal structure of PdCl(L-1). *Inorganic Chemistry Communications.* 7(4) pp. 502-505 (2004)
- [18] Singh, P., & Singh, A. K.: Palladium (II) complexes of tridentate chalcogenated Schiff bases and related ligands of (S, N, S/Se/Te) type: Synthesis and structural chemistry. *Inorganica Chimica Acta.* 387 pp. 441-445 (2012)
- [19] Rooseboom, M., Vermeulen, N. P., Durgut, F., & Commandeur, J. N.: Comparative study on the bioactivation mechanisms and cytotoxicity of Te-phenyl-L-tellurocysteine, Se-phenyl-L-selenocysteine, and S-phenyl-L-cysteine. *Chemical research in toxicology.* 15(12) pp. 1610-1618 (2002)
- [20] Andersson, C. M., Hallberg, A., Brattsand, R., Cotgreave, I. A., Engman, L., & Persson, J.: Glutathione peroxidase-like activity of diaryl tellurides. *Bioorganic & Medicinal Chemistry Letters.* 3(12) pp 2553-2558 (1993)
- [21] Moh. Sattar, Krishna Patidar, Raviraj Ananda Thorat & Sangit Kumar: Copper-Mediated Selective Mono- and Sequential Organochalcogenation of C-H Bonds: Synthesis of Hybrid Unsymmetrical Aryl Ferrocene Chalcogenides. *The Journal of Organic Chemistry.* 84 (11) pp. 6669-6678 (2019)
- [22] Sehnem, J. A., Milani, P., Nascimento, V., Andrade, L. H., Dorneles, L., & Braga, A. L.: Synthesis of new fluororous modular chiral ligand derivatives from amino alcohols and application in enantioselective carbon-carbon bond-forming alkylation reactions. *Tetrahedron: Asymmetry,* 21(8) pp. 997-1003 (2010)

- [23] Vargas, F., Sehnem, J. A., Galetto, F. Z., & Braga, A. L.: Modular chiral β -selenium-, sulfur-, and tellurium amides: Synthesis and application in the palladium-catalyzed asymmetric allylic alkylation. *Tetrahedron*. 64(2) pp. 392-398 (2008)
- [24] Braga, A. L., Paixao, M. W., & Marin, G.: Seleno-imine: a new class of versatile, modular N, Se ligands for asymmetric palladium-catalyzed allylic alkylation. *Synlett*. 2005(11) pp. 1675-1678 (2005)
- [25] Kumar, A., Rao, G. K., Saleem, F., & Singh, A. K.: Organoselenium ligands in catalysis. *Dalton Transactions*. 41(39) pp. 11949-11977 (2012)
- [26] Menon, S. C., Singh, H. B., Patel, R. P., Das, K., & Butcher, R. J.: Synthesis and Reactivity of Chiral Tellurium Azomethines: Pseudopolymorphism of [o-(((1S,2R)-2-Hydroxy-2-phenyl-1-methylethyl)amino) methinyl] phenyl] tellurium(IV) Bromide. *Organometallics*, 16(4) pp. 563-571 (1997)
- [27] Mugesh, G., & Singh, H. B.: Heteroatom-directed aromatic lithiation: A versatile route to the synthesis of organochalcogen (Se, Te) compounds. *Accounts of chemical research*, 35(4) pp. 226-236 (2002)
- [28] Revanna, R. H., Kumar, R. P., Hosamani, A., & Siddagangaiah, P. B.: Efficient and practical synthesis of modular chiral β -organochalcogeno amines, $\text{ArYCH}_2\text{CH(R)NH}_2$, and single crystal structures of (S)- $\text{MsOCH}_2\text{CH(Bz)NH}_3^+\cdot\text{Cl}^-$ and (R)- $\text{MsOCH}_2\text{CH(Ph)NH}_3^+\cdot\text{Cl}^-$. *Journal of Organometallic Chemistry*. 799 pp. 61-69 (2015)
- [29] G.-J. Cheng, X. Zhang, L. W. Chung, L. Xu, Y.-D. Wu: *Computational Organic Chemistry: Bridging Theory and Experiment in Establishing the Mechanisms of Chemical Reactions*. J. Am. Chem. Soc. 137 pp. 1706–1725 (2015)
- [30] W. L. F Armarego, D. D. Perrin: *Purification of laboratory chemicals* fourth edition, Reed Educational and Professional Publishing Ltd., New Delhi (1996)
- [31] K.J. Irgolic, R.A. Zingaro: *Reactions of organotellurium compounds in organometallic synthesis*, in: E. Becker, M. Tsutsui (Eds.), John Wiley and Sons, Inc 2, (1971)
- [32] Zingaro, R. A., Petragnani, N., & Comasseto, J. V.: *Organometallic synthesis*. Vol. 3, 650. Elsevier, New York (1986)
- [33] M. Robb: *New Chemistry with Gaussian 16 & GaussView 6*, (2022). <https://spiral.imperial.ac.uk/handle/10044/1/57650> (accessed June 4, 2023)
- [34] Lu, T., & Chen, F.: Multiwfn: A multifunctional wavefunction analyzer. *Journal of computational chemistry*. 33(5) pp. 580-592 (2012)
- [35] Humphrey, W., Dalke, A., & Schulten, K.: VMD: visual molecular dynamics. *Journal of molecular graphics*. 14(1) pp. 33-38 (1996)
- [36] Rajegowda, H. R., Kumar, P. R., Hosamani, A., & Butcher, R. J.: Synthesis, characterization and determination of absolute structures of palladium complexes of novel chiral acyclic tellurated Schiff base ligands. *New Journal of Chemistry*. 42(8) pp. 6264-6273 (2018)
- [37] H. R. Rajegowda, P. Raghavendra Kumar, & Das, B.: Synthesis, characterization and antioxidant activity of new β -benzylselenated Schiff bases and their palladium complexes. *Journal of Coordination Chemistry*. 75(9–10) pp. 1273–1288 (2022)
- [38] Miar, M., Shiroudi, A., Pourshamsian, K., Oliaey, A. R., & Hatamjafari, F.: Theoretical investigations on the HOMO–LUMO gap and global reactivity descriptor studies, natural bond orbital, and nucleus-independent chemical shifts analyses of 3-phenylbenzo[d]thiazole-2(3H)-imine and its para-substituted derivatives: Solvent and substituent effects. *Journal of Chemical Research*. 45(1-2) pp. 147-158 (2021)