

Antibiotic Activity of Endophytic Bacteria isolated from *Eucheuma cottoni* of North Galesong Sea, Takalar

Hafsan¹, Isna Rasdianah Aziz², Eka Sukmawaty³, St. Aisyah S⁴, Hasyimuddin⁵,
Zulkarnain⁶, Hajrah⁷

¹²³⁴⁵⁶⁷Department of Biology, Faculty of Science and Technology, Universitas Islam Negeri
Alauddin Makassar, Indonesia. 92113

{hafsan.bio@uin-alauddin.ac.id¹, isna-rasdianah@uin-alauddin.ac.id²}

Abstract. *Aeromonas* sp. and *Klebsiella* sp. were two active endophytic microbe isolated from *Eucheuma cottoni* in North Galesong Sea, Takalar district using agar nutrient medium. These isolates then purified by the strike plate on Klinger Iron Agar medium. Fermentation process was carried out using yeast malt broth medium to provide the active compound. The fermented product were supernatant and pellets. The antimicrobial activity was measured using agar diffusion method. The result show that supernatant which only gave activity to *Staphylococcus aureus* and *Escherichia coli* is *Aeromonas* sp.

Keywords: *Aeromonas* sp., Antibiotic, Endophytic bacteria, *Klebsiella* sp., Seaweed.

1 Introduction

Endophytes, microbes which perform ecological function in a plant, complete their life cycle and limit pathogen damage within host plant tissue. Plant-specific or complex interaction with endophytic microbes involves virtually the entire spectrum of mutualistic association, in which plant releases an array of chemical cues as nutrient source for microbe, while microbes transform bioactive compound to promote their host-plant growth and development [1][2][3][4].

Metabolic products, antibiotics, are typically excreted by microbes at limited level that inhibit or kill other microbes. Several genera of *Pseudomonas viridiflava*, *Taxomyces andreanae*, *Streptomyces* sp., *Muscodor* sp., *Aspergillus niger*, *Artemisia annua*, *Neurospora* sp., and *Pestalotiopsis* sp. are well-known endophytic microbes that produce secondary metabolites as antiinsects [5][6], anticancer and antitumour [7][8][9], antifungal [10][9], antiinflammatory [11], antidiabetic [12], antiviral [13], antihelminthic [14], and antimalarial [15][16]. These can be explored prospectively as certain remedies for particular diseases.

Eucheuma cottonii is a widely seaweed cultivated by farmers in North Galesong sea, a sub-district in Takalar district, South Sulawesi. Seaweed becomes a mainstay commodity of Takalar regency, which provides 36.5% of national production, as the third largest national contribution in Indonesia [17][18]. This high production provides an engagement in bioprospecting that endophytic microbes have potential sources of natural product in holding for novel antibiotic discovery. Thus, the synergistic effect against resistant microbe leads to new choices for the treatment of certain diseases. A need for intense and prolonged use of antibiotics in Indonesia,

the discovery and development of new antibiotics remain to address the one important target that ensures new resource for drug discovery.

Based on its potential, it is necessary to evaluate antibiotic activity of endophytic microbes isolated from *Eucheuma cottonii*, especially in North Galesong Sea.

2 Material and Methods

2.1 Pure culture fermentation of endophytic microbe

Pure culture colonies from previous studies (code A1 and A2, not discussed in this study) were taken 1 ose, inoculated in nutrient agar, then incubated at 37°C 24 hours, while on potato dextrose agar, it's incubated at 28°C for 5 days. The suspension was carried out with 2 ml 0.9% NaCl and inoculated in 10 ml of yeast malt broth liquid. 2 ml inoculum was pipetted and put in containing 10 ml of YM-Broth, incubated at 28°C for 24 hours, and shaken at 200 rpm.

2.2 Cell reculture and antibiotic activity test

Staphylococcus aureus and *Escherichia coli* (experimental strains) were taken 1 ose and then inoculated on natrium agar, then incubated at 37°C 24 hours. Microbial suspension was put 1ml into nutrient agar and potato dextrose agar, homogenized and then left 15 min to half solidify. the soaked paper disc is laid with aseptic fermented filtrate, incubated on nutrient agar at 37°C 24 hours, while 28°C for 5 days on potato dextrose agar. The inhibition zone is seen as a clear zone around the disk containing the results of fermentate. The diameter of the clear zone formed was measured and recorded as an active isolate.

2.3 Biochemical activity test

- 2.3.1 Kligers Iron Agar test. Isolate was taken 1 ose, streak to the surface of the agar slant, then incubated at 37°C 24 hours.
- 2.3.2 Motility test. Isolates were taken 1 ose, inoculated vertically until mid-medium, then incubated at 37°C 24 hours.
- 2.3.3 Indol test. Isolates were taken 1 ose, inoculated into medium then homogenized. After incubation at 37°C 24 hours, Ehrlich reagents were added in one drop to the tube.
- 2.3.4 Methyl red test. Isolates were taken 1 ose, inoculated into medium, homogenized, then incubated at 37°C 24 hours, one drop of methyl red indicator were added to the tube.
- 2.3.5 Voges-Proskauer test. Isolates were taken 1 ose, inoculated into medium, homogenized, incubated at 37°C 24 hours, then α -naphthol and KOH were added to the tube.
- 2.3.6 Citrate test. Isolates were taken 1 ose, streak to the agar slant for citrate test, then incubated at 37°C 24 hours
- 2.3.7 Urea test. Isolates were taken 1 ose, streak to the agar slant for urea test, then incubated at 37°C 24 hours
- 2.3.8 Carbohydrate test. This test used several medium: glucose, lactose, sucrose, maltose, mannitol and malonate. Isolates were inoculated to each medium, homogenized, then incubated at 37°C 24 hours.
- 2.3.9 Lysine test. Isolates were taken 1 ose, inoculated into lysine medium, homogenized, then incubated at 37°C 24 hours

2.3.10 Catalase test. Isolates were taken 1 ose and put on clean glass object, then added with 3% H₂O₂ reagent to characterize the bubble formation

3 Results and Discussion

Pure isolates are fermented using yeast malt broth for 24 hours while being shaken with a shaker incubator at 200 rpm, thus separating into two parts. Yeast malt broth is a liquid medium containing yeast extract as a protein source, maltose and dextrose as a carbon source and peptone as an amino acids source, which are needed by bacteria for growth, cell synthesis, energy requirements in metabolism and movement. Physiological salt solution (0.9% NaCl) maintains the osmotic balance and protect the cell wall of bacteria from such osmotic rupture and contraction as ideal ionic strength [19][20]. Shaking process at 200 rpm for 24 hours maintain the microbe life by producing the secondary metabolites as an early stage of growth that may coordinate competitive microbe repression. The antimicrobial substance active against competitive bacteria was called antibiotics, as signaling agents instead of weapons. For this study, antibiotic activity test was conducted to explore the potential of secondary metabolism result in endophytic microbe isolated from *E. cottonii*.

Antimicrobial activity was carried out using *Staphylococcus aureus* representing gram-positive bacteria and *Escherichia coli* as gram-negative bacteria. Selection of both experimental strains due to their pathogenicity through invasiveness, toxins, and antibiotic resistance. *S. aureus* is opportunistic pathogens colonized skin and membranes, then cause a wide range of diseases, including abscesses, pneumonia, meningitis, urinary tract infections, and food poisoning, whereas *E. coli* are the main cause of chronic diarrhea. Isolates that showed activity against microbial were A1, *Aeromonas* sp., while A2 isolates produced were *Kleibsella* sp.




Fig 1. Inhibition zone of isolate A1

Antibiotic activity was carried out by diffusion agar using nutrient agar. Inhibitory zones formed around the paper disk shows that secondary metabolites of isolate A1 have an inhibitory effect on *S. aureus* and *E. coli*. After incubation in day 1 for 24 hours (figure 1), isolate A1 gave inhibition to *S. aureus* with 0.69 cm (average), compared to *E. coli* with 0.68 cm (average). On the second day with incubation period of 48 hours, the diameter of the inhibitory zone was reduced as seen from the clear zone around the paper disk, in *S. aureus* showed 0.64 cm (average), while in *E. coli* was 0.53 cm (average). The third day of 3 x 24 hours incubation, isolates did not inhibit at all, hence there was no clear area around the paper disk as seen in the

S. aureus test with 0.22 cm (average) while in *E. coli*, the inhibition decreases in 0.12 mm (average). Comparison of inhibition zones from the first to the third day has decreased due to optimum point during the incubation period of 24 hours. Cell development was increasing day by day until nutrients depletion in medium, thus bacteria have low ability to inhibit microbes around.

Table 1. Biochemical activity test of endopyhtic microbe

Biochemical activity test	A1	A2
KIA	AL/A -/-	A/A +/-
Motility	-	+
Indol	-	-
Ornitri	-	-
Urea	-	+
Citrate	-	+
MR	-	V
VP	-	-
Glucose	+	+
Lactose	+	+
Sucrose	+	+
Maltosa	+	+
Manitol	+	+
Malonat	-	+
LIA	-	-
3S	v	-
O	-	-
F	-	-
PAD	-	-
Catalase	+	+

KIA test provides that isolate A1 are able to ferment glucose while A2 ferment lactose and sucrose. Isolates showed negative results marked by no black formation around the inoculation area. The black does not form due to the absence of H₂S gas. Motility test showed a positive isolate A2 characterized by spread growth, indicating isolate were motile, while isolate A1 were negative. Urea test showed that the positive A2 isolate was marked by a change in color from red-orange to red-purple, indicating the occurrence of urea hydrolysis. The citrate test showed isolate A1 was negative, whereas in positive A2 isolate with a marked change in green to blue. Microbes were able to use citrate as the only carbon source. MR and VP test for all isolates showed negative results. This test shows whether glucose can be converted to acetone, hence the medium added with alpha-naphthol and KOH reagent, will produce red, the bacteria succesfully produce acetone.

Carbohydrate fermentation test provided positive result of isolate A1 and A2 with gas formation in the durham tube. LIA test is decarboxylase of carboxyl groups from organic molecule where lysine is an amino acid. The acid was produced in the glucose fermentation which reduces the pH of the media. The role of the lysine decarboxylation reactions would favour the amino formation by acid neutralizing, seen in color changes from yellow to purple.

All isolates showed negative result of this test. PAD test also showed negative result of both isolates. After several biochemical tests, it was known that endophytic microbes that could produce antibiotics from *E. cottonii* were *Aeromonas* sp. by showing inhibitory zones in their antibiotic activity tests.

4 Conclusion

Endophytic microbes from marine algae *Eucheuma cottonii* found in North Galesong Sea produce one isolate, *Aeromonas* sp.

References

- [1] P. R. Hardoim *et al.*, “The Hidden World within Plants: Ecological and Evolutionary Considerations for Defining Functioning of Microbial Endophytes,” *Microbiol. Mol. Biol. Rev.*, vol. 79, no. 3, pp. 2261–2268 (2015). doi: 10.1128/MMBR.00050-14
- [2] A. E. Arnold *et al.*, “Fungal endophytes limit pathogen damage in a tropical tree,” *Proc. Natl. Acad. Sci.*, vol. 100, no. 26, pp. 15649–15654 (2003). doi: 10.1073/pnas.2533483100
- [3] D. V. Badri, T. L. Weir, D. Lelie, van der, and J. M. Vivanco, “Rhizosphere chemical dialogues: plant–microbe interactions,” *Curr. Opin. Biotechnol.*, vol. 20, no. 6, pp. 642–650 (2013). doi: 10.1016/j.copbio.2009.09.014
- [4] G. A. Strobel, “Endophytes as sources of bioactive products,” *Microbes Infect.*, vol. 5, no. 6, pp. 535–544 (2003). doi: 10.1016/S1286-4579(03)00073-X
- [5] Y. J. Cardoza, K. D. Klepzig, and K. F. Raffa, “Bacteria in oral secretions of an endophytic insect inhibit antagonistic fungi,” *Ecol. Entomol.*, vol. 31, no. 6, pp. 636–645 (2006). doi: 10.1111/j.1365-2311.2006.00829.x
- [6] M. C. González, A. L. Anaya, A. E. Glenn, M. L. Macías-Rubalcava, B. E. Hernández-Bautista, and R. T. Hanlin, “*Muscodor yucatanensis*, a new endophytic ascomycete from Mexican chakah, *Bursera simaruba*,” *Mycotaxon*, vol. 110, no. 1, pp. 363–372 (2009).
- [7] J. Li *et al.*, “Antitumor and antimicrobial activities of endophytic streptomycetes from pharmaceutical plants in rainforest,” *Lett. Appl. Microbiol.*, vol. 47, no. 6, pp. 574–580 (2008). doi: 10.1111/j.1472-765X.2008.02470.x
- [8] R. K. Tenguria and F. N. Khan, “Distribution of Endophytic Fungi in Leaves of *Azadirachta indica* A. JUSS. (Neem) of Panchmarhi Biosphere Reserve,” *Curr. Bot.*, vol. 2, no. 2, pp. 27–29 (2011).
- [9] Y. Huang, J. Wang, G. Li, Z. Zheng, and W. Su, “Antitumor and antifungal activities in endophytic fungi isolated from pharmaceutical plants *Taxus mairei*, *Cephalataxus fortunei* and *Torreya grandis*,” *FEMS Immunol. Med. Microbiol.*, vol. 31, no. 2, pp. 163–167 (2001). doi: 10.1111/j.1574-695X.2001.tb00513.x
- [10] C. H. Liu, W. X. Zou, H. Lu, and R. X. Tan, “Antifungal activity of *Artemisia annua* endophyte cultures against phytopathogenic fungi,” *J. Biotechnol.*, vol. 88, no. 3, pp. 277–282 (2001). doi: 10.1016/S0168-1656(01)00285-1
- [11] T. Taechowisan, A. Wanbanjob, P. Tuntiwachwuttikul, and J. Liu, “Anti-inflammatory activity of lansais from endophytic *Streptomyces* sp. SUC1 in LPS-induced RAW 264.7 cells,” *Food Agric. Immunol.*, vol. 20, no. 1, pp. 67–77 (2009). doi: 10.1080/09540100902730064
- [12] V. J. Akshatha, M. S. Nalini, C. D’Souza, and H. S. Prakash, “Streptomycete endophytes from anti-diabetic medicinal plants of the Western Ghats inhibit alpha-amylase and promote glucose uptake,” *Lett. Appl. Microbiol.*, vol. 58, no. 5, pp. 433–439 (2014). doi:

10.1111/lam.12209

- [13] P. Wang *et al.*, “Diketopiperazine Derivatives from the Marine-Derived Actinomycete *Streptomyces* sp. FXJ7.328,” *Mar. drug*, vol. 11, no. 4, pp. 1035–1049 (2013). doi: 10.3390/md11041035
- [14] S. Ōmura *et al.*, “An anthelmintic compound, nafuredin, shows selective inhibition of complex I in helminth mitochondria,” *Proc. Natl. Acad. Sci.*, vol. 98, no. 1, pp. 60–62 (2001). doi: 10.1073/pnas.98.1.60
- [15] M. S. Baba *et al.*, “In vivo antimalarial activity of the endophytic actinobacteria, *Streptomyces* SUK 10,” *Microb. Physiol. Biochem.*, vol. 53, no. 12, pp. 847–855 (2015). doi: 10.1007/s12275-015-5076-6
- [16] S. Wiyakrutta *et al.*, “Endophytic fungi with anti-microbial, anti-cancer and anti-malarial activities isolated from Thai medicinal plants,” *World J. Microbiol. Biotechnol.*, vol. 20, no. 3, pp. 265–272 (2004). doi: 10.1023/B:WIBI.0000023832.27679.a8
- [17] A. Rahadiati, K. Soewardi, Y. Wardiatno, and D. Sutrisno, “Pemetaan sebaran budidaya rumput laut: pendekatan analisis multitemporal (studi kasus di kabupaten Takalar Sulawesi Selatan),” *Maj. Ilm. Globe*, vol. 20, no. 1, pp. 13–22 (2018).
- [18] K. Yaqin, I. Burhanuddin, and W. Samad, “Biodiversity of seaweed and their metal contents from littoral zone of South Sulawesi waters,” *Semin. Int. Kelautan, Univ. Udayana Bali*, pp. 1–16 (2011).
- [19] D. K. Maheshwari, *Bacterial Diversity in Sustainable Agriculture*. Switzerland: Springer, Cham, (2014). doi: 10.1007/978-3-319-05936-5
- [20] R. D. Sleator and C. Hill, “Bacterial osmoadaptation: the role of osmolytes in bacterial stress and virulence,” *Microbiol. Rev.*, vol. 26, no. 1, pp. 49–71 (2002). doi: 10.1111/j.1574-6976.2002.tb00598.x